

Human Haptoglobin (Hpt) ELISA Kit | Hpt elisa kit

Human Haptoglobin (Hpt) ELISA Kit

CATALOG #
AAA63966

REACTIVITY
Human


PRODUCT SYNONYMS

Haptoglobin (Hpt); Human Haptoglobin (Hpt) ELISA Kit; HP Hp2-Alpha Alpha-2-Macroglobulin Zonulin; Hpt elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of HIF2a. No significant cross-reactivity or interference between HIF2a and analogues was observed.
Sequence Length	347
Samples	Mouse Serum, Plasma, Tissue Homogenates, Cell Lysates, Cell Culture Supernates Or Other Biological Fluids
Assay Type	Quantitative Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.053ng/mL.
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level HIF2a were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level HIF2a were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for Hpt elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of HIF2a in mouse serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to HIF2a. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to HIF2a. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain HIF2a, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of HIF2a in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	186910296
NCBI GeneID	3240
NCBI Accession #	NP_001119574.1
NCBI GenBank Nucleotide #	NM_001126102.2
UniProt Accession #	P00738; P00737; Q0VAC4; Q0VAC5; Q2PP15; Q3B7J0; Q6LBY9; Q9UC67; B0AZL5
Molecular Weight	38,452 Da
NCBI Official Full Name	haptoglobin isoform 2 preproprotein
NCBI Official Synonym Full Names	haptoglobin
NCBI Official Symbol	HP
NCBI Official Synonym Symbols	BP; HPA1S; HP2ALPHA2
NCBI Protein Information	haptoglobin
UniProt Protein Name	Haptoglobin
UniProt Gene Name	HP

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.