

Mouse Androgen Receptor (AR) ELISA Kit | AR elisa kit

Mouse Androgen Receptor (AR) ELISA Kit

CATALOG #
AAA64060

REACTIVITY
Mouse


PRODUCT SYNONYMS

Androgen Receptor (AR); Mouse Androgen Receptor (AR) ELISA Kit; AIS; DHTR; HUMARA; KD; NR3C4; SBMA; SMAX1; TFM; Dihydrotestosterone receptor; Nuclear Receptor Subfamily 3; Group C; Member 4; Testicular Feminization; Kennedy Disease; AR elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of AR. No significant cross-reactivity or interference between AR and analogues was observed.
Sequence Length	899
Samples	Tissue homogenates, cell lysates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.061ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level AR were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level AR were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for AR elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of AR in mouse tissue homogenates, cell lysates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to AR. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to AR. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain AR, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of AR in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	7304901
NCBI GeneID	11835
NCBI Accession #	NP_038504.1
NCBI GenBank Nucleotide #	NM_013476.4
UniProt Accession #	P19091
Molecular Weight	98,194 Da
NCBI Official Full Name	androgen receptor
NCBI Official Synonym Full Names	androgen receptor
NCBI Official Symbol	Ar
NCBI Official Synonym Symbols	Tfm; AW320017
NCBI Protein Information	androgen receptor
UniProt Protein Name	Androgen receptor
UniProt Gene Name	Ar
UniProt Synonym Gene Names	Nr3c4

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-elisa-kits/androgen-receptor-ar/64060>