

Mouse Intercellular Adhesion Molecule 1 (ICAM1) ELISA Kit | ICAM1 elisa kit

Mouse Intercellular Adhesion Molecule 1 (ICAM1) ELISA Kit

CATALOG #
AAA64201

REACTIVITY
Mouse


PRODUCT SYNONYMS

Intercellular Adhesion Molecule 1 (ICAM1); Mouse Intercellular Adhesion Molecule 1 (ICAM1) ELISA Kit; PAPP-A; PAPA; ASBAP2; DIPLA1; IGFBP-4ase; PAPPA1; Pappalysin 1;; ICAM1 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of ICAM1. No significant cross-reactivity or interference between ICAM1 and analogues was observed.
Sequence Length	537
Assay Type	Quantitative Sandwich
Samples	Mouse serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids
Detection Range	0.312-20ng/mL
Sensitivity	< 0.134ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level ICAM1 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level ICAM1 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for ICAM1 elisa kit	Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to ICAM1. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to ICAM1. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain ICAM1, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of ICAM1 in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of ICAM1 in mouse serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	21389311
NCBI GeneID	15894
NCBI Accession #	NP_034623.1
NCBI GenBank Nucleotide #	NM_010493.3
UniProt Accession #	P13597; Q61828
Molecular Weight	58,753 Da
NCBI Official Full Name	intercellular adhesion molecule 1
NCBI Official Synonym Full Names	intercellular adhesion molecule 1
NCBI Official Symbol	Icam1
NCBI Official Synonym Symbols	CD54; Ly-47; Icam-1; MALA-2
NCBI Protein Information	intercellular adhesion molecule 1
UniProt Protein Name	Intercellular adhesion molecule 1
UniProt Gene Name	Icam1
UniProt Synonym Gene Names	Icam-1; ICAM-1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-elisa-kits/intercellular-adhesion-molecule-1-icam1/64201>