

Human Kynurenine-3-Monooxygenase (KMO) ELISA Kit | KMO elisa kit

Human Kynurenine-3-Monooxygenase (KMO) ELISA Kit

CATALOG #
AAA64248

REACTIVITY
Human


PRODUCT SYNONYMS

Kynurenine-3-Monooxygenase (KMO); Human Kynurenine-3-Monooxygenase (KMO) ELISA Kit; Kynurenine 3-Hydroxylase; KMO elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of KMO. No significant cross-reactivity or interference between KMO and analogues was observed.
Sequence Length	486
Samples	Tissue homogenates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.781-50ng/mL
Sensitivity	0.24ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level KMO were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level KMO were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for KMO elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of KMO in human tissue homogenates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to KMO. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to KMO. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain KMO, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of KMO in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	187960045
NCBI GeneID	8564
NCBI Accession #	NP_003670.2
NCBI GenBank Nucleotide #	NM_003679.4
UniProt Accession #	O15229; Q5SY07; Q5SY08; Q5SY09; A2A2U8; A2A2U9; A2A2V0
Molecular Weight	51,682 Da
NCBI Official Full Name	kynurenine 3-monooxygenase
NCBI Official Synonym Full Names	kynurenine 3-monooxygenase
NCBI Official Symbol	KMO
NCBI Official Synonym Symbols	dJ317G22.1
NCBI Protein Information	kynurenine 3-monooxygenase
UniProt Protein Name	Kynurenine 3-monooxygenase
UniProt Gene Name	KMO

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.