

Human Neutral Sphingomyelinase (NSMASE) ELISA Kit | NSMASE elisa kit

Human Neutral Sphingomyelinase (NSMASE) ELISA Kit

CATALOG #
AAA64254

REACTIVITY
Human


PRODUCT SYNONYMS

Neutral Sphingomyelinase (NSMASE); Human Neutral Sphingomyelinase (NSMASE) ELISA Kit; SMPD2; N-SMase; ISC1; Neutral Sphingomyelin Phosphodiesterase; Sphingomyelin Phosphodiesterase 2; Neutral Membrane; Lyso-platelet-activating factor-phospholipase C; NSMASE elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of NSMASE. No significant cross-reactivity or interference between NSMASE and analogues was observed.
Sequence Length	423
Samples	Tissue homogenates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.067ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level NSMASE were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level NSMASE were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for NSMASE elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of NSMASE in human tissue homogenates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to NSMASE. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to NSMASE. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain NSMASE, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of NSMASE in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	92859618
NCBI GeneID	6610
NCBI Accession #	NP_003071.2
NCBI GenBank Nucleotide #	NM_003080.2
UniProt Accession #	O60906; Q5TED1; Q9BWR3
Molecular Weight	47,646 Da
NCBI Official Full Name	sphingomyelin phosphodiesterase 2
NCBI Official Synonym Full Names	sphingomyelin phosphodiesterase 2
NCBI Official Symbol	SMPD2
NCBI Official Synonym Symbols	ISC1; NSMASE; NSMASE1
NCBI Protein Information	sphingomyelin phosphodiesterase 2
UniProt Protein Name	Sphingomyelin phosphodiesterase 2
UniProt Gene Name	SMPD2
UniProt Synonym Gene Names	Lyso-PAF-PLC; N-SMase; nSMase

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/neutral-sphingomyelinase-nsmase/64254>