

Mouse C Reactive Protein (CRP) ELISA Kit | CRP elisa kit

Mouse C Reactive Protein (CRP) ELISA Kit

CATALOG #
AAA64349

REACTIVITY
Mouse


PRODUCT SYNONYMS

C Reactive Protein (CRP); Mouse C Reactive Protein (CRP) ELISA Kit; C-RP; PTX1; Pentraxin-Related; Pentraxin 1; CRP elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of CRP. No significant cross-reactivity or interference between CRP and analogues was observed.
Sequence Length	225
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.073ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level CRP were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level CRP were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CRP elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of CRP in mouse serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to CRP. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to CRP. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain CRP, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of CRP in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	162138926
NCBI GeneID	12944
NCBI Accession #	NP_031794.3
NCBI GenBank Nucleotide #	NM_007768.4
UniProt Accession #	P14847; E9PZ20
Molecular Weight	25,360 Da
NCBI Official Full Name	C-reactive protein
NCBI Official Synonym Full Names	C-reactive protein, pentraxin-related
NCBI Official Symbol	Crp
NCBI Official Synonym Symbols	AI255847
NCBI Protein Information	C-reactive protein
UniProt Protein Name	C-reactive protein
UniProt Gene Name	Crp
UniProt Synonym Gene Names	Ptx1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.