

Mouse N-Acetyltransferase 1 (NAT1) ELISA Kit | NAT1 elisa kit

Mouse N-Acetyltransferase 1 (NAT1) ELISA Kit

CATALOG #
AAA64387

REACTIVITY
Mouse


PRODUCT SYNONYMS

N-Acetyltransferase 1 (NAT1); Mouse N-Acetyltransferase 1 (NAT1) ELISA Kit; AAC1; Arylamine N-Acetyltransferase; Arylamide acetylase 1; Monomorphic arylamine N-acetyltransferase; NAT1 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of NAT1. No significant cross-reactivity or interference between NAT1 and analogues was observed.
Sequence Length	868
Samples	Tissue homogenates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.066ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level NAT1 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level NAT1 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for NAT1 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of NAT1 in mouse tissue homogenates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to NAT1. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to NAT1. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain NAT1, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of NAT1 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	110630015
NCBI GeneID	13690
NCBI Accession #	NP_001035221.1
NCBI GenBank Nucleotide #	NM_001040131.2
UniProt Accession #	Q62448; P97867; Q0VGZ3; Q3UNC1; Q5XKD8; Q6P1Z0; Q921U3
Molecular Weight	97,894 Da
NCBI Official Full Name	eukaryotic translation initiation factor 4 gamma 2 isoform 2
NCBI Official Synonym Full Names	eukaryotic translation initiation factor 4, gamma 2
NCBI Official Symbol	Eif4g2
NCBI Official Synonym Symbols	p97; Nat1; DAP-5; Natm1; E130105L11Rik
NCBI Protein Information	eukaryotic translation initiation factor 4 gamma 2
UniProt Protein Name	Eukaryotic translation initiation factor 4 gamma 2
UniProt Gene Name	Eif4g2
UniProt Synonym Gene Names	eIF-4-gamma 2; eIF-4G 2; eIF4G 2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.