

Human Syndecan 3 (SDC3) ELISA Kit | SDC3 elisa kit

Human Syndecan 3 (SDC3) ELISA Kit

CATALOG #
AAA64557

REACTIVITY
Human


PRODUCT SYNONYMS

Syndecan 3 (SDC3); Human Syndecan 3 (SDC3) ELISA Kit; SDCN SYND3 N-Syndecan Syndecan Proteoglycan 3; SDC3 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of SDC3. No significant cross-reactivity or interference between SDC3 and analogues was observed.
Sequence Length	442
Samples	Tissue homogenates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.625-40ng/mL
Sensitivity	0.226ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level SDC3 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level SDC3 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for SDC3 elisa kit

Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of SDC3 in human tissue homogenates and other biological fluids.

Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to SDC3. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to SDC3. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain SDC3, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of SDC3 in the samples is then determined by comparing the O.D. of the samples to the standard curve.

NCBI AND UNIPROT INFORMATION

NCBI GI #	187960059
NCBI GeneID	9672
NCBI Accession #	NP_055469.3
NCBI GenBank Nucleotide #	NM_014654.3
UniProt Accession #	O75056; Q5T1Z6; Q5T1Z7; Q96CT3; Q96PR8
Molecular Weight	45,497 Da
NCBI Official Full Name	syndecan-3
NCBI Official Synonym Full Names	syndecan 3
NCBI Official Symbol	SDC3
NCBI Official Synonym Symbols	SDCN; SYND3
NCBI Protein Information	syndecan-3
UniProt Protein Name	Syndecan-3
UniProt Gene Name	SDC3
UniProt Synonym Gene Names	KIAA0468; SYND3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/syndecan-3-sdc3/64557>