

Human Histone Deacetylase 1 (HDAC1) ELISA Kit | HDAC1 elisa kit

Human Histone Deacetylase 1 (HDAC1) ELISA Kit

CATALOG #
AAA64567

REACTIVITY
Human


PRODUCT SYNONYMS

Histone Deacetylase 1 (HDAC1); Human Histone Deacetylase 1 (HDAC1) ELISA Kit; HDAC1; GON10; HD1; RPD3; RPD3L1; HDAC1 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of HDAC1. No significant cross-reactivity or interference between HDAC1 and analogues was observed.
Sequence Length	482
Samples	Tissue homogenates, cell lysates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.061ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level HDAC1 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level HDAC1 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for HDAC1 elisa kit	Intended Uses: This kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of HDAC1 in human tissue homogenates, cell lysates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to HDAC1. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to HDAC1. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain HDAC1, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of HDAC1 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	13128860
NCBI GeneID	3065
NCBI Accession #	NP_004955.2
NCBI GenBank Nucleotide #	NM_004964.2
UniProt Accession #	Q13547; Q92534
Molecular Weight	55,103 Da
NCBI Official Full Name	histone deacetylase 1
NCBI Official Synonym Full Names	histone deacetylase 1
NCBI Official Symbol	HDAC1
NCBI Official Synonym Symbols	HD1; RPD3; GON-10; RPD3L1
NCBI Protein Information	histone deacetylase 1
UniProt Protein Name	Histone deacetylase 1
UniProt Gene Name	HDAC1
UniProt Synonym Gene Names	RPD3L1; HD1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/histone-deacetylase-1-hdac1/64567>