

Mouse Mannose Receptor C Type 1 (MRC1) ELISA Kit | MRC1 elisa kit

Mouse Mannose Receptor C Type 1 (MRC1) ELISA Kit

CATALOG #
AAA64669

REACTIVITY
Mouse


PRODUCT SYNONYMS

Mannose Receptor C Type 1 (MRC1); Mouse Mannose Receptor C Type 1 (MRC1) ELISA Kit; CD206; CLEC13D; CLEC13DL; MRC1L1; C-type lectin domain family 13 member D-like; Macrophage mannose receptor 1-like protein 1; MRC1 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of MRC1. No significant cross-reactivity or interference between MRC1 and analogues was observed.
Sequence Length	1456
Samples	Serum, plasma, tissue homogenates, cell lysates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.059ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level MRC1 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level MRC1 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for MRC1 elisa kit	Intended Uses: This kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of MRC1 in mouse serum, plasma, tissue homogenates, cell lysates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to MRC1. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to MRC1. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain MRC1, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of MRC1 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	224967062
NCBI GeneID	17533
NCBI Accession #	NP_032651.2
NCBI GenBank Nucleotide #	NM_008625.2
UniProt Accession #	Q61830; Q8C502
Molecular Weight	164,981 Da
NCBI Official Full Name	macrophage mannose receptor 1
NCBI Official Synonym Full Names	mannose receptor, C type 1
NCBI Official Symbol	Mrc1
NCBI Official Synonym Symbols	MR; CD206; AW259686
NCBI Protein Information	macrophage mannose receptor 1
UniProt Protein Name	Macrophage mannose receptor 1
UniProt Gene Name	Mrc1
UniProt Synonym Gene Names	MMR

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-elisa-kits/mannose-receptor-c-type-1-mrc1/64669>