

Mouse Sirtuin 1 (SIRT1) ELISA Kit | SIRT1 elisa kit

Mouse Sirtuin 1 (SIRT1) ELISA Kit

CATALOG #
AAA64726

REACTIVITY
Mouse


PRODUCT SYNONYMS

Sirtuin 1 (SIRT1); Mouse Sirtuin 1 (SIRT1) ELISA Kit; SIR2L1; Silent Mating Type Information Regulation 2 Homolog 1; NAD-Dependent Deacetylase Sirtuin-1; SIR2-like protein 1; SirtT1 75 kDa fragment; SIRT1 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of SIRT1. No significant cross-reactivity or interference between SIRT1 and analogues was observed.
Sequence Length	737
Assay Type	Quantitative Sandwich
Samples	Mouse serum, plasma, tissue homogenates, cell lysates or other biological fluids
Detection Range	0.156-10ng/mL
Sensitivity	< 0.055ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level SIRT1 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level SIRT1 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for SIRT1 elisa kit	Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to SIRT1. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to SIRT1. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain SIRT1, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of SIRT1 in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of SIRT1 in mouse serum, plasma, tissue homogenates, cell lysates or other biological fluids.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	9790229
NCBI GeneID	93759
NCBI Accession #	NP_062786.1
NCBI GenBank Nucleotide #	NM_019812.3
UniProt Accession #	Q923E4; Q9QXG8
Molecular Weight	59,875 Da
NCBI Official Full Name	NAD-dependent protein deacetylase sirtuin-1 isoform 1
NCBI Official Synonym Full Names	sirtuin 1
NCBI Official Symbol	Sirt1
NCBI Official Synonym Symbols	Sir2; Sir2a; SIR2L1; AA673258; Sir2alpha
NCBI Protein Information	NAD-dependent protein deacetylase sirtuin-1
UniProt Protein Name	NAD-dependent protein deacetylase sirtuin-1
UniProt Gene Name	Sirt1
UniProt Synonym Gene Names	Sir2l1; Sir2; mSIR2a; 75Sir1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.