

Mouse Cytochrome C, Somatic (CYCS) ELISA Kit | CYCS elisa kit

Mouse Cytochrome C, Somatic (CYCS) ELISA Kit

CATALOG #
AAA64877

REACTIVITY
Mouse


PRODUCT SYNONYMS

Cytochrome C; Somatic (CYCS); Mouse Cytochrome C; Somatic (CYCS) ELISA Kit; Cyt-C; HCS; CYC; CytC; CYCS elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of CYCS. No significant cross-reactivity or interference between CYCS and analogues was observed.
Sequence Length	105
Samples	Tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.058ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level CYCS were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level CYCS were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CYCS elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of CYCS in mouse tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to CYCS. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to CYCS. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain CYCS, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of CYCS in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	6681095
NCBI GeneID	13063
NCBI Accession #	NP_031834.1
NCBI GenBank Nucleotide #	NM_007808.4
UniProt Accession #	P62897; P00009; Q8C2S2
Molecular Weight	11,605 Da
NCBI Official Full Name	cytochrome c, somatic
NCBI Official Synonym Full Names	cytochrome c, somatic
NCBI Official Symbol	Cycs
NCBI Protein Information	cytochrome c, somatic
UniProt Protein Name	Cytochrome c, somatic
UniProt Gene Name	Cycs

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.