

Mouse Betacellulin (bTC) ELISA Kit | bTC elisa kit

Mouse Betacellulin (bTC) ELISA Kit

CATALOG #
AAA64962

REACTIVITY
Mouse


PRODUCT SYNONYMS

Betacellulin (bTC); Mouse Betacellulin (bTC) ELISA Kit; Probetacellulin; bTC elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of BTC. No significant cross-reactivity or interference between BTC and analogues was observed.
Sequence Length	177
Samples	Serum, plasma, tissue homogenates, cell lysates and other biological fluids.
Assay Type	Sandwich
Detection Range	15.625-1000pg/mL
Sensitivity	7.3pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level BTC were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level BTC were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for bTC elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of bTC in mouse serum, plasma, tissue homogenates, cell lysates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to BTC. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to BTC. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain BTC, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of BTC in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	6680812
NCBI GeneID	12223
NCBI Accession #	NP_031594.1
NCBI GenBank Nucleotide #	NM_007568.5
UniProt Accession #	Q543J8; Q05928
Molecular Weight	19,664 Da
NCBI Official Full Name	betacellulin preproprotein
NCBI Official Synonym Full Names	betacellulin, epidermal growth factor family member
NCBI Official Symbol	Btc
NCBI Official Synonym Symbols	Bcn
NCBI Protein Information	betacellulin
UniProt Protein Name	Betacellulin
UniProt Gene Name	Btc

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.