

Human Sirtuin 4 (SIRT4) ELISA Kit | SIRT4 elisa kit

Human Sirtuin 4 (SIRT4) ELISA Kit

CATALOG #
AAA64965

REACTIVITY
Human


PRODUCT SYNONYMS

Sirtuin 4 (SIRT4); Human Sirtuin 4 (SIRT4) ELISA Kit; SIR2L4; Silent Mating Type Information Regulation 2 Homolog 4; SIR2-like protein 4; NAD-dependent protein deacetylase sirtuin-4; SIRT4 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of SIRT4. No significant cross-reactivity or interference between SIRT4 and analogues was observed.
Sequence Length	314
Assay Type	Quantitative Sandwich
Samples	Human serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids
Detection Range	0.156-10ng/mL
Sensitivity	< 0.051ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level SIRT4 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level SIRT4 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for SIRT4 elisa kit	Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to SIRT4. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to SIRT4. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain SIRT4, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of SIRT4 in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of SIRT4 in human serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	6912662
NCBI GeneID	23409
NCBI Accession #	NP_036372.1
NCBI GenBank Nucleotide #	NM_012240.2
UniProt Accession #	Q9Y6E7; O43346; Q32M33
Molecular Weight	35,188 Da
NCBI Official Full Name	NAD-dependent protein lipoamidase sirtuin-4, mitochondrial
NCBI Official Synonym Full Names	sirtuin 4
NCBI Official Symbol	SIRT4
NCBI Official Synonym Symbols	SIR2L4
NCBI Protein Information	NAD-dependent protein lipoamidase sirtuin-4, mitochondrial
UniProt Protein Name	NAD-dependent protein lipoamidase sirtuin-4, mitochondrial
UniProt Gene Name	SIRT4

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.