

Mouse Activated Leukocyte Cell Adhesion Molecule (ALCAM) ELISA Kit | ALCAM elisa kit

Mouse Activated Leukocyte Cell Adhesion Molecule (ALCAM) ELISA Kit

CATALOG #
AAA64997

REACTIVITY
Mouse


PRODUCT SYNONYMS

Activated Leukocyte Cell Adhesion Molecule (ALCAM); Mouse Activated Leukocyte Cell Adhesion Molecule (ALCAM) ELISA Kit; CD166; MEMD; Activated Leucocyte Cell Adhesion Molecule; ALCAM elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of ALCAM. No significant cross-reactivity or interference between ALCAM and analogues was observed.
Sequence Length	583
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.468-30ng/mL
Sensitivity	0.13ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level ALCAM were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level ALCAM were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for ALCAM elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of ALCAM in mouse serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to ALCAM. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to ALCAM. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain ALCAM, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of ALCAM in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	31791059
NCBI GeneID	11658
NCBI Accession #	NP_033785.1
NCBI GenBank Nucleotide #	NM_009655.3
UniProt Accession #	Q61490; O70136; Q8CDA5; Q8R2T0
Molecular Weight	65,092 Da
NCBI Official Full Name	CD166 antigen isoform 1
NCBI Official Synonym Full Names	activated leukocyte cell adhesion molecule
NCBI Official Symbol	Alcam
NCBI Official Synonym Symbols	BEN; SC1; MuSC; CD166; AI853494; DM-GRASP
NCBI Protein Information	CD166 antigen
UniProt Protein Name	CD166 antigen
UniProt Gene Name	Alcam

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-elisa-kits/activated-leukocyte-cell-adhesion-molecule-alcam/64997>