

Human Triggering Receptor Expressed On Myeloid Cells 2 (TREM2) ELISA Kit | TREM2 elisa kit

Human Triggering Receptor Expressed On Myeloid Cells 2 (TREM2) ELISA Kit

CATALOG # AAA65041	REACTIVITY Human
------------------------------	----------------------------


PRODUCT SYNONYMS

Triggering Receptor Expressed On Myeloid Cells 2 (TREM2); Human Triggering Receptor Expressed On Myeloid Cells 2 (TREM2) ELISA Kit; TREM2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of TREM2. No significant cross-reactivity or interference between TREM2 and analogues was observed.
Sequence Length	219
Assay Type	Quantitative Sandwich
Samples	Human serum, plasma, tissue homogenates, cell lysates or other biological fluids
Detection Range	62.5-4000pg/mL
Sensitivity	< 21.9pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level TREM2 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level TREM2 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for TREM2 elisa kit	Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to TREM2. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to TREM2. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain TREM2, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of TREM2 in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of TREM2 in human serum, plasma, tissue homogenates, cell lysates or other biological fluids.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	425876764
NCBI GeneID	54209
NCBI Accession #	NP_001258750.1
NCBI GenBank Nucleotide #	NM_001271821.1
UniProt Accession #	Q9NZC2; Q8N5H8; Q8WYN6
Molecular Weight	24,669 Da
NCBI Official Full Name	triggering receptor expressed on myeloid cells 2 isoform 2
NCBI Official Synonym Full Names	triggering receptor expressed on myeloid cells 2
NCBI Official Symbol	TREM2
NCBI Official Synonym Symbols	TREM-2; Trem2a; Trem2b; Trem2c
NCBI Protein Information	triggering receptor expressed on myeloid cells 2
UniProt Protein Name	Triggering receptor expressed on myeloid cells 2
UniProt Gene Name	TREM2
UniProt Synonym Gene Names	TREM-2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/triggering-receptor-expressed-on-myeloid-cells-2-trem2/65041>