

Human Low Density Lipoprotein Receptor (LDLR) ELISA Kit | LDLR elisa kit

Human Low Density Lipoprotein Receptor (LDLR) ELISA Kit

CATALOG #
AAA65054

REACTIVITY
Human


PRODUCT SYNONYMS

Low Density Lipoprotein Receptor (LDLR); Human Low Density Lipoprotein Receptor (LDLR) ELISA Kit; HMG-R; HMGY; LDLR elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of LDLR. No significant cross-reactivity or interference between LDLR and analogues was observed.
Sequence Length	860
Assay Type	Quantitative Sandwich
Samples	Human tissue homogenates, cell lysates or other biological fluids
Detection Range	0.312-20ng/mL
Sensitivity	< 0.118ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level LDLR were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level LDLR were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for LDLR elisa kit	Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to LDLR. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to LDLR. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain LDLR, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of LDLR in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of LDLR in human tissue homogenates, cell lysates or other biological fluids.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504975
NCBI GeneID	3949
NCBI Accession #	NP_000518.1
NCBI GenBank Nucleotide #	NM_000527.4
UniProt Accession #	P01130; Q53ZD9; Q59FQ1; Q9UDH7; B4DII3; B4DJZ8; B4DR00; B4DTQ3; C0JYY8; H0YLU8; H0YNT7
Molecular Weight	82,255 Da
NCBI Official Full Name	low-density lipoprotein receptor isoform 1
NCBI Official Synonym Full Names	low density lipoprotein receptor
NCBI Official Symbol	LDLR
NCBI Official Synonym Symbols	FH; FHC; LDLCQ2
NCBI Protein Information	low-density lipoprotein receptor
UniProt Protein Name	Low-density lipoprotein receptor
UniProt Gene Name	LDLR
UniProt Synonym Gene Names	LDL receptor

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/low-density-lipoprotein-receptor-ldlr/65054>