

Rat Transgelin (TAGLN) ELISA Kit | TAGLN elisa kit

Rat Transgelin (TAGLN) ELISA Kit

CATALOG # AAA65056	REACTIVITY Rat
------------------------------	--------------------------


PRODUCT SYNONYMS

Transgelin (TAGLN); Rat Transgelin (TAGLN) ELISA Kit; SMCC; SM22; TAGLN1; WS3-10; Transgelin Variant 2; 22 kDa actin-binding protein; Protein WS3-10; Smooth muscle protein 22-alpha; TAGLN elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of TAGLN. No significant cross-reactivity or interference between TAGLN and analogues was observed.
Sequence Length	201
Samples	Serum, plasma, tissue homogenates and other biological fluids.
Assay Type	Sandwich
Detection Range	1.56-100ng/mL
Sensitivity	0.58ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level TAGLN were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level TAGLN were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for TAGLN elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of TAGLN in rat serum, plasma, tissue homogenates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to TAGLN. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to TAGLN. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain TAGLN, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of TAGLN in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	13928744
NCBI GeneID	25123
NCBI Accession #	NP_113737.1
NCBI GenBank Nucleotide #	NM_031549.2
UniProt Accession #	P31232
Molecular Weight	22,603 Da
NCBI Official Full Name	transgelin
NCBI Official Synonym Full Names	transgelin
NCBI Official Symbol	Tagln
NCBI Official Synonym Symbols	Sm22
NCBI Protein Information	transgelin
UniProt Protein Name	Transgelin
UniProt Gene Name	Tagln
UniProt Synonym Gene Names	Sm22; SM22-alpha

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.