

Human Sirtuin 6 (SIRT6) ELISA Kit | SIRT6 elisa kit

Human Sirtuin 6 (SIRT6) ELISA Kit

CATALOG #
AAA65132

REACTIVITY
Human


PRODUCT SYNONYMS

Sirtuin 6 (SIRT6); Human Sirtuin 6 (SIRT6) ELISA Kit; SIR2L6; Silent Mating Type Information Regulation 2 Homolog 6; Mono-ADP-Ribosyltransferase Sirtuin-6; Regulatory protein SIR2 homolog 6; SIR2-like protein 6; SIRT6 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of SIRT6. No significant cross-reactivity or interference between SIRT6 and analogues was observed.
Sequence Length	328
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.312-20ng/mL
Sensitivity	0.124ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level SIRT6 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level SIRT6 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for SIRT6 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of SIRT6 in human serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to SIRT6. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to SIRT6. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain SIRT6, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of SIRT6 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	300797597
NCBI GeneID	51548
NCBI Accession #	NP_001180214.1
NCBI GenBank Nucleotide #	NM_001193285.2
UniProt Accession #	Q8N6T7; O75291; Q6IAF5; Q6PK99; Q8NCD2; Q9BSI5; Q9BWP3; Q9NRC7; Q9UQD1; B2RCD0
Molecular Weight	36,065 Da
NCBI Official Full Name	NAD-dependent protein deacetylase sirtuin-6 isoform 2
NCBI Official Synonym Full Names	sirtuin 6
NCBI Official Symbol	SIRT6
NCBI Official Synonym Symbols	SIR2L6
NCBI Protein Information	NAD-dependent protein deacetylase sirtuin-6
UniProt Protein Name	NAD-dependent protein deacetylase sirtuin-6
UniProt Gene Name	SIRT6
UniProt Synonym Gene Names	SIR2L6

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.