

Mouse Replication Initiator 1 (REPIN1) ELISA Kit | REPIN1 elisa kit

Mouse Replication Initiator 1 (REPIN1) ELISA Kit

CATALOG #
AAA65157

REACTIVITY
Mouse


PRODUCT SYNONYMS

Replication Initiator 1 (REPIN1); Mouse Replication Initiator 1 (REPIN1) ELISA Kit; REPIN1 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of REPIN1. No significant cross-reactivity or interference between REPIN1 and analogues was observed.
Sequence Length	598
Samples	Tissue homogenates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.312-20ng/mL
Sensitivity	0.135ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level REPIN1 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level REPIN1 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for REPIN1 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of REPIN1 in mouse tissue homogenates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to REPIN1. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to REPIN1. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain REPIN1, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of REPIN1 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	347658943
NCBI GeneID	58887
NCBI Accession #	NP_001073370.2
NCBI GenBank Nucleotide #	NM_001079901.1
UniProt Accession #	Q5U4E2; Q8BTQ5
Molecular Weight	61,830 Da
NCBI Official Full Name	replication initiator 1 isoform a
NCBI Official Synonym Full Names	replication initiator 1
NCBI Official Symbol	Repin1
NCBI Official Synonym Symbols	Ap4; Zfp464; Zfp-464; AI425994; E430037F08Rik
NCBI Protein Information	replication initiator 1
UniProt Protein Name	Replication initiator 1
UniProt Gene Name	Repin1
UniProt Synonym Gene Names	Zfp464; Zfp-464

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.