

Mouse Versican (VS) ELISA Kit | VS elisa kit

Mouse Versican (VS) ELISA Kit

CATALOG # AAA65171	REACTIVITY Mouse
------------------------------	----------------------------


PRODUCT SYNONYMS

Versican (VS); Mouse Versican (VS) ELISA Kit; CSPG2; VCAN; ERVR; PG-M; WGN1; Chondroitin Sulfate Proteoglycan 2; Chondroitin sulfate proteoglycan core protein 2; Glial hyaluronate-binding; Large fibroblast proteoglycan; VS elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of VS. No significant cross-reactivity or interference between VS and analogues was observed.
Sequence Length	3357
Samples	Serum, plasma, tissue homogenates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.059ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level VS were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level VS were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for VS elisa kit	Intended Uses: This kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of VS in mouse serum, plasma, tissue homogenates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to VS. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to VS. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain VS, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of VS in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	108935831
NCBI GeneID	13003
UniProt Accession #	Q62059; Q62058; Q9CUU0
Molecular Weight	74,177 Da
NCBI Official Full Name	Versican core protein
NCBI Official Synonym Full Names	versican
NCBI Official Symbol	Vcan
NCBI Official Synonym Symbols	NG2; hdf; PG-M; Cspg2; DPEAAE; PG-M(V0); PG-M(V1); 9430051N09; 5430420N07Rik
NCBI Protein Information	versican core protein
UniProt Protein Name	Versican core protein
UniProt Gene Name	Vcan
UniProt Synonym Gene Names	Cspg2; Chondroitin sulfate proteoglycan 2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.