

Human Cadherin 17 (CDH17) ELISA Kit | CDH17 elisa kit

Human Cadherin 17 (CDH17) ELISA Kit

CATALOG #
AAA65234

REACTIVITY
Human


PRODUCT SYNONYMS

Cadherin 17 (CDH17); Human Cadherin 17 (CDH17) ELISA Kit; CDH16; HPT-1; HPT1; LI-Cadherin; Cadherin; Liver Intestine; Intestinal peptide-associated transporter HPT-1; Liver-intestine cadherin; CDH17 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of CDH17. No significant cross-reactivity or interference between CDH17 and analogues was observed.
Sequence Length	832
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.054ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level CDH17 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level CDH17 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CDH17 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of CDH17 in human serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to CDH17. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to CDH17. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain CDH17, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of CDH17 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	221316595
NCBI GeneID	1015
NCBI Accession #	NP_001138135.1
NCBI GenBank Nucleotide #	NM_001144663.1
UniProt Accession #	Q12864; Q15336; Q2M2E0
Molecular Weight	92,219 Da
NCBI Official Full Name	cadherin-17
NCBI Official Synonym Full Names	cadherin 17
NCBI Official Symbol	CDH17
NCBI Official Synonym Symbols	HPT1; CDH16; HPT-1
NCBI Protein Information	cadherin-17
UniProt Protein Name	Cadherin-17
UniProt Gene Name	CDH17
UniProt Synonym Gene Names	LI-cadherin

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.