

Human Hexokinase 2 (HK2) ELISA Kit | HK2 elisa kit

Human Hexokinase 2 (HK2) ELISA Kit

CATALOG #
AAA65251

REACTIVITY
Human


PRODUCT SYNONYMS

Hexokinase 2 (HK2); Human Hexokinase 2 (HK2) ELISA Kit; HK II; Muscle form hexokinase; HK2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of HK2. No significant cross-reactivity or interference between HK2 and analogues was observed.
Sequence Length	917
Samples	Serum, plasma, tissue homogenates or other biological fluids
Assay Type	Quantitative Sandwich
Detection Range	31.2-2000pg/mL
Sensitivity	< 12.9pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level HK2 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level HK2 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for HK2 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of HK2 in human serum, plasma, tissue homogenates or other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to HK2. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to HK2. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain HK2, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of HK2 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	15553127
NCBI GeneID	3099
NCBI Accession #	NP_000180.2
NCBI GenBank Nucleotide #	NM_000189.4
UniProt Accession #	P52789; Q8WU87; Q9UN82; D6W5J2
Molecular Weight	102,380 Da
NCBI Official Full Name	hexokinase-2
NCBI Official Synonym Full Names	hexokinase 2
NCBI Official Symbol	HK2
NCBI Official Synonym Symbols	HKII; HXK2
NCBI Protein Information	hexokinase-2
UniProt Protein Name	Hexokinase-2
UniProt Gene Name	HK2
UniProt Synonym Gene Names	HK II

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.