

Rabbit Superoxide Dismutase 3, Extracellular (SOD3) ELISA Kit | SOD3 elisa kit

Rabbit Superoxide Dismutase 3, Extracellular (SOD3) ELISA Kit

CATALOG #
AAA65314

REACTIVITY
Rabbit


PRODUCT SYNONYMS

Superoxide Dismutase 3; Extracellular (SOD3); Rabbit Superoxide Dismutase 3; Extracellular (SOD3) ELISA Kit; SOD-3; EC-SOD; Extracellular Superoxide Dismutase Cu-Zn; SOD3 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rabbit
Specificity	This assay has high sensitivity and excellent specificity for detection of SOD3. No significant cross-reactivity or interference between SOD3 and analogues was observed.
Sequence Length	244
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.781-50ng/mL
Sensitivity	0.28ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level SOD3 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level SOD3 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for SOD3 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of SOD3 in rabbit serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to SOD3. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to SOD3. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain SOD3, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of SOD3 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	130508879
NCBI GeneID	100009320
NCBI Accession #	NP_001076101.1
NCBI GenBank Nucleotide #	NM_001082632.1
UniProt Accession #	P41975
Molecular Weight	25,688 Da
NCBI Official Full Name	extracellular superoxide dismutase
NCBI Official Symbol	SOD3
NCBI Protein Information	extracellular superoxide dismutase [Cu-Zn]
UniProt Protein Name	Extracellular superoxide dismutase [Cu-Zn]
UniProt Gene Name	SOD3
UniProt Synonym Gene Names	EC-SOD

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.