

Human Peroxisome Proliferator Activated Receptor Gamma (PPARg) ELISA Kit | PPARg elisa kit

Human Peroxisome Proliferator Activated Receptor Gamma (PPARg) ELISA Kit

CATALOG # AAA65481	REACTIVITY Human
------------------------------	----------------------------


PRODUCT SYNONYMS

Peroxisome Proliferator Activated Receptor Gamma (PPARg); Human Peroxisome Proliferator Activated Receptor Gamma (PPARg) ELISA Kit; PPAR-G; PPARG1; PPARG2; NR1C3; Glitazone Receptor; Nuclear Receptor Subfamily 1 Group C Member 3; PPARg elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of PPARG. No significant cross-reactivity or interference between PPARG and analogues was observed.
Sequence Length	477
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.781-50ng/mL
Sensitivity	0.29ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level PPARG were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level PPARG were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for PPARG elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of PPARG in human serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to PPARG. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to PPARG. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain PPARG, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of PPARG in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	116284368
NCBI GeneID	5468
NCBI Accession #	NP_005028.4
NCBI GenBank Nucleotide #	NM_005037.5
UniProt Accession #	P37231; O00684; O00710; O14515; Q0QJH8; Q15178; Q15179; Q15180; Q15832; Q86U60; A8K3G6; B5BUA1
Molecular Weight	21,580 Da
NCBI Official Full Name	peroxisome proliferator-activated receptor gamma isoform 1
NCBI Official Synonym Full Names	peroxisome proliferator activated receptor gamma
NCBI Official Symbol	PPARG
NCBI Official Synonym Symbols	GLM1; CIMT1; NR1C3; PPARG1; PPARG2; PPARGgamma
NCBI Protein Information	peroxisome proliferator-activated receptor gamma
UniProt Protein Name	Peroxisome proliferator-activated receptor gamma
UniProt Gene Name	PPARG
UniProt Synonym Gene Names	NR1C3; PPAR-gamma

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/peroxisome-proliferator-activated-receptor-gamma-pparg/65481>