

Rat Apolipoprotein C2 (APOC2) ELISA Kit | APOC2 elisa kit

Rat Apolipoprotein C2 (APOC2) ELISA Kit

CATALOG #
AAA65521

REACTIVITY
Rat


PRODUCT SYNONYMS

Apolipoprotein C2 (APOC2); Rat Apolipoprotein C2 (APOC2) ELISA Kit; APC2; Apo-C2; ProapoC-II; Apolipoprotein C-II; Proapolipoprotein C-II; APOC2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of APOC2. No significant cross-reactivity or interference between APOC2 and analogues was observed.
Sequence Length	97
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	39.062-2500pg/mL
Sensitivity	14pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level APOC2 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level APOC2 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for APOC2 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of APOC2 in rat serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to APOC2. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to APOC2. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain APOC2 biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of APOC2 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	145553986
NCBI GeneID	292697
NCBI Accession #	NP_001078821.1
NCBI GenBank Nucleotide #	NM_001085352.1
UniProt Accession #	G3V8D4
Molecular Weight	10,695 Da
NCBI Official Full Name	apolipoprotein C-II
NCBI Official Synonym Full Names	apolipoprotein C2
NCBI Official Symbol	Apoc2
NCBI Official Synonym Symbols	RGD1560725
NCBI Protein Information	apolipoprotein C-II
UniProt Protein Name	Apolipoprotein C-II (Predicted)
UniProt Gene Name	Apoc2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.