

Mouse B-Cell Activating Factor (BAFF) ELISA Kit | BAFF elisa kit

Mouse B-Cell Activating Factor (BAFF) ELISA Kit

CATALOG #
AAA65553

REACTIVITY
Mouse


PRODUCT SYNONYMS

B-Cell Activating Factor (BAFF); Mouse B-Cell Activating Factor (BAFF) ELISA Kit; CD257; TNFSF13B; BLYS; TALL1; THANK; TNFSF20; ZTNF4; B lymphocyte stimulator; Tumor Necrosis Factor Ligand Superfamily Member 13B; Dendritic cell-derived TNF-like; BAFF elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of BAFF. No significant cross-reactivity or interference between BAFF and analogues was observed.
Sequence Length	290
Samples	Serum, plasma, tissue homogenates and other biological fluids.
Assay Type	Sandwich
Detection Range	1.56-100pg/mL
Sensitivity	0.55pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level BAFF were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level BAFF were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for BAFF elisa kit	Intended Uses: This kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of BAFF in mouse serum, plasma, tissue homogenates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to BAFF. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to BAFF. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain BAFF, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of BAFF in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	15826858
NCBI GeneID	24099
NCBI Accession #	NP_296371.1
NCBI GenBank Nucleotide #	NM_033622.1
UniProt Accession #	Q9WU72; Q7TQ58
Molecular Weight	32,166 Da
NCBI Official Full Name	tumor necrosis factor ligand superfamily member 13B isoform 1
NCBI Official Synonym Full Names	tumor necrosis factor (ligand) superfamily, member 13b
NCBI Official Symbol	Tnfsf13b
NCBI Official Synonym Symbols	BAFF; BLYS; TALL1; THANK; zTNF4; TALL-1; Tnlg7a; TNFSF20; D8ErtD387e
NCBI Protein Information	tumor necrosis factor ligand superfamily member 13B
UniProt Protein Name	Tumor necrosis factor ligand superfamily member 13B
UniProt Gene Name	Tnfsf13b
UniProt Synonym Gene Names	Baff

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.