

Horse Interleukin 10 (IL10) ELISA Kit | IL10 elisa kit

Horse Interleukin 10 (IL10) ELISA Kit

CATALOG #
AAA65615

REACTIVITY
Horse


PRODUCT SYNONYMS

Interleukin 10 (IL10); Horse Interleukin 10 (IL10) ELISA Kit; AR; CRDGF; SDGF; Colorectum Cell-Derived Growth Factor; Schwannoma-Derived Growth Factor; IL10 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Horse
Specificity	This assay has high sensitivity and excellent specificity for detection of IL10. No significant cross-reactivity or interference between IL10 and analogues was observed.
Sequence Length	178
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	15.625-1000pg/mL
Sensitivity	5.4pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level IL10 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level IL10 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for IL10 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of IL10 in equine serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to IL10. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to IL10. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain IL10, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of IL10 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	126723435
NCBI GeneID	100034187
NCBI Accession #	NP_001075959.1
NCBI GenBank Nucleotide #	NM_001082490.1
UniProt Accession #	Q28374
Molecular Weight	20,451 Da
NCBI Official Full Name	interleukin-10
NCBI Official Symbol	IL10
NCBI Official Synonym Symbols	IL-10
NCBI Protein Information	interleukin-10
UniProt Protein Name	Interleukin-10
UniProt Gene Name	IL10
UniProt Synonym Gene Names	IL-10; CSIF

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.