

Rat Neurochondrin (NCDN) ELISA Kit | NCDN elisa kit

Rat Neurochondrin (NCDN) ELISA Kit

CATALOG #
AAA65685

REACTIVITY
Rat


PRODUCT SYNONYMS

Neurochondrin (NCDN); Rat Neurochondrin (NCDN) ELISA Kit; Norbin; NCDN elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of NCDN. No significant cross-reactivity or interference between NCDN and analogues was observed.
Sequence Length	729
Samples	Tissue homogenates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.055ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level NCDN were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level NCDN were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for NCDN elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of NCDN in rat tissue homogenates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to NCDN. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to NCDN. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain NCDN, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of NCDN in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	139947663
NCBI GeneID	89791
NCBI Accession #	NP_445995.2
NCBI GenBank Nucleotide #	NM_053543.2
UniProt Accession #	O35095; Q5PQW2
Molecular Weight	78,923 Da
NCBI Official Full Name	neurochondrin
NCBI Official Synonym Full Names	neurochondrin
NCBI Official Symbol	Ncdn
NCBI Official Synonym Symbols	Norbin
NCBI Protein Information	neurochondrin
UniProt Protein Name	Neurochondrin
UniProt Gene Name	Ncdn

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.