

Rabbit Heparin Cofactor II (HCII) ELISA Kit | HCII elisa kit

Rabbit Heparin Cofactor II (HCII) ELISA Kit

CATALOG #
AAA65741

REACTIVITY
Rabbit


PRODUCT SYNONYMS

Heparin Cofactor II (HCII); Rabbit Heparin Cofactor II (HCII) ELISA Kit; SERPIND1; HC2; HCF2; HC II; HLS2; LS2; Serpin D1; Serpin Peptidase Inhibitor Clade D Member 1; Protease inhibitor leuserpin-2; HCII elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rabbit
Specificity	This assay has high sensitivity and excellent specificity for detection of HCII. No significant cross-reactivity or interference between HCII and analogues was observed.
Sequence Length	480
Samples	Plasma.
Assay Type	Sandwich
Detection Range	3.125-200U/L
Sensitivity	1.03U/L
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level HCII were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level HCII were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for HCII elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of HCII in rabbit plasma. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to HCII. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to HCII. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain HCII, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of HCII in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	126722597
NCBI GeneID	100009170
NCBI Accession #	NP_001075798.1
NCBI GenBank Nucleotide #	NM_001082329.1
UniProt Accession #	P47776
Molecular Weight	54,532 Da
NCBI Official Full Name	heparin cofactor 2
NCBI Official Symbol	SERPIND1
NCBI Official Synonym Symbols	HCII
NCBI Protein Information	heparin cofactor 2
UniProt Protein Name	Heparin cofactor 2
UniProt Gene Name	SERPIND1
UniProt Synonym Gene Names	HCF2; HC-II

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/rabbit-elisa-kits/heparin-cofactor-ii-hcii/65741>