

Rat Dopamine Transporter (DAT) ELISA Kit | DAT elisa kit

Rat Dopamine Transporter (DAT) ELISA Kit

CATALOG #
AAA65836

REACTIVITY
Rat


PRODUCT SYNONYMS

Dopamine Transporter (DAT); Rat Dopamine Transporter (DAT) ELISA Kit; DAT elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of DAT. No significant cross-reactivity or interference between DAT and analogues was observed.
Sequence Length	619
Samples	Tissue homogenates, cell lysates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.312-20ng/mL
Sensitivity	0.142ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level DAT were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level DAT were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for DAT elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of DAT in Rat tissue homogenates, cell lysates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to DAT. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to DAT. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain DAT, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of DAT in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	7106413
NCBI GeneID	24898
NCBI Accession #	NP_036826.1
NCBI GenBank Nucleotide #	NM_012694.2
UniProt Accession #	P23977
Molecular Weight	68,746 Da
NCBI Official Full Name	sodium-dependent dopamine transporter
NCBI Official Synonym Full Names	solute carrier family 6 member 3
NCBI Official Symbol	Slc6a3
NCBI Official Synonym Symbols	Dat1
NCBI Protein Information	sodium-dependent dopamine transporter
UniProt Protein Name	Sodium-dependent dopamine transporter
UniProt Gene Name	Slc6a3
UniProt Synonym Gene Names	DA transporter; DAT

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/rat-elisa-kits/dopamine-transporter-dat/65836>