

Human Cystatin 3 (CST3) ELISA Kit | CST3 elisa kit

Human Cystatin 3 (CST3) ELISA Kit

 CATALOG #
AAA65865

 REACTIVITY
Human


PRODUCT SYNONYMS

Cystatin 3 (CST3); Human Cystatin 3 (CST3) ELISA Kit; Cys-C Cystatin C Gamma Trace Post-Gamma-Globulin Neuroendocrine Basic Polypeptide Amyloid Angiopathy And Cerebral Hemorrhage; CST3 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of CST3. No significant cross-reactivity or interference between CST3 and analogues was observed.
Sequence Length	146
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	1.56-100ng/mL
Sensitivity	0.64ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level CST3 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level CST3 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CST3 elisa kit

Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of CST3 in human serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.

Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to CST3. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to CST3. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain CST3, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of CST3 in the samples is then determined by comparing the O.D. of the samples to the standard curve.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4503107
NCBI GeneID	1471
NCBI Accession #	NP_000090.1
NCBI GenBank Nucleotide #	NM_000099.3
UniProt Accession #	P01034; Q6FGW9; B2R5J9; D3DW42
Molecular Weight	15,799 Da
NCBI Official Full Name	cystatin-C
NCBI Official Synonym Full Names	cystatin C
NCBI Official Symbol	CST3
NCBI Official Synonym Symbols	ARMD11; HEL-S-2
NCBI Protein Information	cystatin-C
UniProt Protein Name	Cystatin-C
UniProt Gene Name	CST3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/cystatin-3-cst3/65865>