

Mouse Fibroblast Growth Factor 12 (FGF12) ELISA Kit | FGF12 elisa kit

Mouse Fibroblast Growth Factor 12 (FGF12) ELISA Kit

CATALOG #
AAA65887

REACTIVITY
Mouse


PRODUCT SYNONYMS

Fibroblast Growth Factor 12 (FGF12); Mouse Fibroblast Growth Factor 12 (FGF12) ELISA Kit; FGF12B; FHF1; Fibroblast Growth Factor Homologous Factor 1; Myocyte-Activating Factor; FGF12 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of FGF12. No significant cross-reactivity or interference between FGF12 and analogues was observed.
Sequence Length	207
Samples	Serum, plasma and other biological fluids.
Assay Type	Sandwich
Detection Range	15.625-1000pg/mL
Sensitivity	6.3pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level FGF12 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level FGF12 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for FGF12 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of FGF12 in mouse serum, plasma and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to FGF12. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to FGF12. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain FGF12, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of FGF12 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	449310780
NCBI GeneID	14167
NCBI Accession #	NP_001263348.1
NCBI GenBank Nucleotide #	NM_001276419.2
UniProt Accession #	P61329; O35339; P70376; Q924B4; Q92912; Q93001
Molecular Weight	20,424 Da
NCBI Official Full Name	fibroblast growth factor 12 isoform c
NCBI Official Synonym Full Names	fibroblast growth factor 12
NCBI Official Symbol	Fgf12
NCBI Official Synonym Symbols	Fhf1; FHF-1; FGF-12; AV114868; B230343J05Rik
NCBI Protein Information	fibroblast growth factor 12
UniProt Protein Name	Fibroblast growth factor 12
UniProt Gene Name	Fgf12
UniProt Synonym Gene Names	Fhf1; FGF-12; FHF-1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.