

Human Chemokine C-X-C-Motif Ligand 16 (CXCL16) ELISA Kit | CXCL16 elisa kit

Human Chemokine C-X-C-Motif Ligand 16 (CXCL16) ELISA Kit

CATALOG #
AAA66127

REACTIVITY
Human


PRODUCT SYNONYMS

Chemokine C-X-C-Motif Ligand 16 (CXCL16); Human Chemokine C-X-C-Motif Ligand 16 (CXCL16) ELISA Kit; SR-PSOX; CXCLG16; SRPSOX; Scavenger receptor for phosphatidylserine and oxidized low density lipoprotein; Small-inducible cytokine B16; CXCL16 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of CXCL16. No significant cross-reactivity or interference between CXCL16 and analogues was observed.
Sequence Length	273
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.067ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level CXCL16 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level CXCL16 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CXCL16 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of CXCL16 in human serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to CXCL16. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to CXCL16. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain CXCL16, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of CXCL16 in the samples is then determined by comparing the O.D. of the samples to the standard curve. CALCULATI
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	154816178
NCBI GeneID	58191
NCBI Accession #	NP_001094282.1
NCBI GenBank Nucleotide #	NM_001100812.1
UniProt Accession #	Q9H2A7; Q8TC80; Q96K63; Q9BXD6; Q9H2F6; A0N0N4; A8K7U9; B2RCB0
Molecular Weight	27,579 Da
NCBI Official Full Name	C-X-C motif chemokine 16
NCBI Official Synonym Full Names	C-X-C motif chemokine ligand 16
NCBI Official Symbol	CXCL16
NCBI Official Synonym Symbols	SRPSOX; CXCLG16; SR-PSOX
NCBI Protein Information	C-X-C motif chemokine 16
UniProt Protein Name	C-X-C motif chemokine 16
UniProt Gene Name	CXCL16
UniProt Synonym Gene Names	SCYB16; SRPSOX; SR-PSOX

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.