

Human Statherin (STATH) ELISA Kit | STATH elisa kit

Human Statherin (STATH) ELISA Kit

CATALOG #
AAA66175

REACTIVITY
Human


PRODUCT SYNONYMS

Statherin (STATH); Human Statherin (STATH) ELISA Kit; STR; STATH elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of STATH. No significant cross-reactivity or interference between STATH and analogues was observed.
Sequence Length	52
Samples	Serum, plasma, tissue homogenates and other biological fluids.
Assay Type	Sandwich
Detection Range	3.125-200ng/mL
Sensitivity	1.33ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level STATH were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level STATH were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for STATH elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of STATH in human serum, plasma, tissue homogenates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to STATH. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to STATH. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain STATH, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of STATH in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	57164946
NCBI GeneID	6779
NCBI Accession #	NP_001009181.1
NCBI GenBank Nucleotide #	NM_001009181.1
UniProt Accession #	P02808; A6NKE9; B2R4F8
Molecular Weight	6,073 Da
NCBI Official Full Name	statherin isoform b
NCBI Official Synonym Full Names	statherin
NCBI Official Symbol	STATH
NCBI Official Synonym Symbols	STR
NCBI Protein Information	statherin
UniProt Protein Name	Statherin
UniProt Gene Name	STATH

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/statherin-stath/66175>