

Mouse DNA Repair Protein RAD50 (RAD50) ELISA Kit | RAD50 elisa kit

Mouse DNA Repair Protein RAD50 (RAD50) ELISA Kit

CATALOG # AAA66180	REACTIVITY Mouse
------------------------------	----------------------------


PRODUCT SYNONYMS

DNA Repair Protein RAD50 (RAD50); Mouse DNA Repair Protein RAD50 (RAD50) ELISA Kit; RAD50 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of RAD50. No significant cross-reactivity or interference between RAD50 and analogues was observed.
Sequence Length	1312
Samples	Tissue homogenates, cell lysates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.312-20ng/mL
Sensitivity	0.113ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, qmiddle and high level RAD50 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level RAD50 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for RAD50 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of RAD50 in mouse tissue homogenates, cell lysates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to RAD50. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to RAD50. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain RAD50, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of RAD50 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	60392985
NCBI GeneID	19360
UniProt Accession #	P70388; Q6PEB0; Q8C2T7; Q9CU59
Molecular Weight	57,589 Da
NCBI Official Full Name	DNA repair protein RAD50
NCBI Official Synonym Full Names	RAD50 double strand break repair protein
NCBI Official Symbol	Rad50
NCBI Official Synonym Symbols	Mrell; Rad50l
NCBI Protein Information	DNA repair protein RAD50
UniProt Protein Name	DNA repair protein RAD50
UniProt Gene Name	Rad50
UniProt Synonym Gene Names	mRad50

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-elisa-kits/dna-repair-protein-rad50-rad50/66180>