

Human Fibroblast Growth Factor 17 (FGF17) ELISA Kit | FGF17 elisa kit

Human Fibroblast Growth Factor 17 (FGF17) ELISA Kit

CATALOG #
AAA66363

REACTIVITY
Human


PRODUCT SYNONYMS

Fibroblast Growth Factor 17 (FGF17); Human Fibroblast Growth Factor 17 (FGF17) ELISA Kit; FGF17 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of FGF17. No significant cross-reactivity or interference between FGF17 and analogues was observed.
Sequence Length	205
Samples	Serum, plasma, tissue homogenates, cell lysates and other biological fluids.
Assay Type	Sandwich
Detection Range	78.125-5000pg/mL
Sensitivity	35.96pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level FGF17 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level FGF17 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for FGF17 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of FGF17 in human serum, plasma, tissue homogenates, cell lysates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to FGF17. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to FGF17. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain FGF17, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of FGF17 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	751247036
NCBI GeneID	8822
NCBI Accession #	NP_001291407.1
NCBI GenBank Nucleotide #	NM_001304478.1
UniProt Accession #	O60258; Q2M2W1; B7ZLG4
Molecular Weight	23,669 Da
NCBI Official Full Name	fibroblast growth factor 17 isoform 2
NCBI Official Synonym Full Names	fibroblast growth factor 17
NCBI Official Symbol	FGF17
NCBI Official Synonym Symbols	HH20; FGF-13; FGF-17
NCBI Protein Information	fibroblast growth factor 17
UniProt Protein Name	Fibroblast growth factor 17
UniProt Gene Name	FGF17
UniProt Synonym Gene Names	FGF-17

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.