

Human CD200 Receptor 2 (CD200R2) ELISA Kit | CD200R2 elisa kit

Human CD200 Receptor 2 (CD200R2) ELISA Kit

CATALOG #
AAA66371

REACTIVITY
Human


PRODUCT SYNONYMS

CD200 Receptor 2 (CD200R2); Human CD200 Receptor 2 (CD200R2) ELISA Kit; CD200R1L; CD200RLa; HuCD200R2; Cell surface glycoprotein CD200 receptor 2; CD200 cell surface glycoprotein receptor-like a; Cell surface glycoprotein OX2 receptor 2; CD200R2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of CD200R2. No significant cross-reactivity or interference between CD200R2 and analogues was observed.
Sequence Length	271
Samples	Serum, plasma, tissue homogenates, cell lysates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.781-50ng/mL
Sensitivity	0.34ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level CD200R2 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level CD200R2 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CD200R2 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of CD200R2 in human serum, plasma, tissue homogenates, cell lysates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to CD200R2. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to CD200R2. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain CD200R2, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of CD200R2 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	151301013
NCBI GeneID	344807
NCBI Accession #	NP_001008784.2
NCBI GenBank Nucleotide #	NM_001008784.2
UniProt Accession #	Q6Q8B3; Q6WHB7
Molecular Weight	27,771 Da
NCBI Official Full Name	cell surface glycoprotein CD200 receptor 2 isoform 1
NCBI Official Synonym Full Names	CD200 receptor 1 like
NCBI Official Symbol	CD200R1L
NCBI Official Synonym Symbols	CD200R2; CD200RLa
NCBI Protein Information	cell surface glycoprotein CD200 receptor 2
UniProt Protein Name	Cell surface glycoprotein CD200 receptor 2
UniProt Gene Name	CD200R1L
UniProt Synonym Gene Names	CD200R2; CD200 receptor-like 2; HuCD200R2; CD200RLa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.