

Mouse Proenkephalin (PENK) ELISA Kit | PENK elisa kit

Mouse Proenkephalin (PENK) ELISA Kit

CATALOG #
AAA66397

REACTIVITY
Mouse


PRODUCT SYNONYMS

Proenkephalin (PENK); Mouse Proenkephalin (PENK) ELISA Kit; Preproenkephalin; Proenkephalin-A; Synenkephalin; Met-enkephalin; Met-enkephalin-Arg-Gly-Leu; Leu-enkephalin; Met-enkephalin-Arg-Phe; PENK elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of PENK. No significant cross-reactivity or interference between PENK and analogues was observed.
Sequence Length	268
Assay Type	Quantitative Sandwich
Samples	Mouse serum, plasma, tissue homogenates or other biological fluids
Detection Range	78.1-5000pg/mL
Sensitivity	< 21.3pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level PENK were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level PENK were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for PENK elisa kit	Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to PENK. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to PENK. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain PENK, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of PENK in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of PENK in mouse serum, plasma, tissue homogenates or other biological fluids.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	50950119
NCBI GeneID	18619
NCBI Accession #	NP_001002927.1
NCBI GenBank Nucleotide #	NM_001002927.3
UniProt Accession #	P22005; Q68G73
Molecular Weight	31,004 Da
NCBI Official Full Name	proenkephalin-A
NCBI Official Synonym Full Names	preproenkephalin
NCBI Official Symbol	Penk
NCBI Official Synonym Symbols	ENK; PPA; Penk1; AI326464
NCBI Protein Information	proenkephalin-A
UniProt Protein Name	Proenkephalin-A
UniProt Gene Name	Penk
UniProt Synonym Gene Names	Penk1; OGF

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.