

Rat Interleukin 17F (IL17F) ELISA Kit | IL17F elisa kit

Rat Interleukin 17F (IL17F) ELISA Kit

CATALOG # AAA66570	REACTIVITY Rat
------------------------------	--------------------------


PRODUCT SYNONYMS

Interleukin 17F (IL17F); Rat Interleukin 17F (IL17F) ELISA Kit; ML-1; IL17F elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of IL17F. No significant cross-reactivity or interference between IL17F and analogues was observed.
Sequence Length	161
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	15.625-1000pg/mL
Sensitivity	5.7pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level IL17F were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level IL17F were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for IL17F elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of IL17F in mouse serum, plasma, Other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to IL17F. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to IL17F. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain IL17F, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of IL17F in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	845634515
NCBI GeneID	301291
NCBI Accession #	NP_001015011.2
NCBI GenBank Nucleotide #	NM_001015011.2
UniProt Accession #	Q5BJ95
Molecular Weight	17,712 Da
NCBI Official Full Name	interleukin-17F
NCBI Official Synonym Full Names	interleukin 17F
NCBI Official Symbol	IL17f
NCBI Official Synonym Symbols	IL-17F
NCBI Protein Information	interleukin-17F
UniProt Protein Name	Interleukin-17F
UniProt Gene Name	IL17f
UniProt Synonym Gene Names	IL-17F

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.