

Mouse Calreticulin (CRT) ELISA Kit | CRT elisa kit

Mouse Calreticulin (CRT) ELISA Kit

CATALOG #
AAA66640

REACTIVITY
Mouse


PRODUCT SYNONYMS

Calreticulin (CRT); Mouse Calreticulin (CRT) ELISA Kit; CALR; RO; SSA; CC1qR; ERp60; HACBP; grp60; CRTc; CRP55; Calregulin; Sicca Syndrome Antigen A; Autoantigen Ro; Endoplasmic reticulum resident protein 60; CRT elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of CRT. No significant cross-reactivity or interference between CRT and analogues was observed.
Sequence Length	416
Samples	Serum, plasma, tissue homogenates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.067ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level CRT were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level CRT were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CRT elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of CRT in mouse serum, plasma, tissue homogenates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to CRT. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to CRT. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain CRT, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of CRT in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	6680836
NCBI GeneID	12317
NCBI Accession #	NP_031617.1
NCBI GenBank Nucleotide #	NM_007591.3
UniProt Accession #	P14211; Q3TVD2
Molecular Weight	47,995 Da
NCBI Official Full Name	calreticulin
NCBI Official Synonym Full Names	calreticulin
NCBI Official Symbol	Calr
NCBI Official Synonym Symbols	CRT; Calregulin
NCBI Protein Information	calreticulin
UniProt Protein Name	Calreticulin
UniProt Gene Name	Calr
UniProt Synonym Gene Names	ERp60

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-elisa-kits/calreticulin-crt/66640>