

Human Synuclein Alpha (SNCa) ELISA Kit | SNCa elisa kit

Human Synuclein Alpha (SNCa) ELISA Kit

CATALOG #
AAA66649

REACTIVITY
Human


PRODUCT SYNONYMS

Synuclein Alpha (SNCa); Human Synuclein Alpha (SNCa) ELISA Kit; SNC-A; aSYN; PD1; PARK1; PARK4; NACP; Non-A Beta Component Of Alzheimer's Disease Amyloid Precursor Protein; Non-A4 component of amyloid precursor; SNCa elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of SNCa. No significant cross-reactivity or interference between SNCa and analogues was observed.
Sequence Length	140
Samples	Serum, plasma or other biological fluids
Assay Type	Quantitative Sandwich
Detection Range	15.6-1000pg/mL
Sensitivity	< 5.9pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level SNCa were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level SNCa were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for SNCA elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of SNCA in human serum, plasma or other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to SNCA. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to SNCA. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain SNCA, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of SNCA in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507109
NCBI GeneID	6622
NCBI Accession #	NP_000336.1
NCBI GenBank Nucleotide #	NM_000345.3
UniProt Accession #	P37840; Q13701; Q4JHI3; Q6IAU6; A8K2A4
Molecular Weight	13,109 Da
NCBI Official Full Name	alpha-synuclein isoform NACP140
NCBI Official Synonym Full Names	synuclein alpha
NCBI Official Symbol	SNCA
NCBI Official Synonym Symbols	PD1; NACP; PARK1; PARK4
NCBI Protein Information	alpha-synuclein
UniProt Protein Name	Alpha-synuclein
UniProt Gene Name	SNCA
UniProt Synonym Gene Names	NACP; PARK1; NACP

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.