

Mouse N-Acetyl Beta-D-Glucosaminidase (NAGase) ELISA Kit | NAGase elisa kit

Mouse N-Acetyl Beta-D-Glucosaminidase (NAGase) ELISA Kit

CATALOG #
AAA66678

REACTIVITY
Mouse


PRODUCT SYNONYMS

N-Acetyl Beta-D-Glucosaminidase (NAGase); Mouse N-Acetyl Beta-D-Glucosaminidase (NAGase) ELISA Kit; MGEA5 MEA5 NCOAT Hexosaminidase C N-Acetylhexosaminidase Hyaluronidase Meningioma-Expressed Antigen 5 Nuclear Cytoplasmic O-GlcNAcase And Acetyltransferase; NAGase elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of NAGase. No significant cross-reactivity or interference between NAGase and analogues was observed.
Sequence Length	916
Samples	Urine.
Assay Type	Sandwich
Detection Range	0.781-50ng/mL
Sensitivity	0.50ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level NAGase were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level NAGase were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for NAGase elisa kit

Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of NAGase in mouse urine.

Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to NAGase. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to NAGase. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain NAGase, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of NAGase in the samples is then determined by comparing the O.D. of the samples to the standard curve.

NCBI AND UNIPROT INFORMATION

NCBI GI #	15011884
NCBI GeneID	76055
NCBI Accession #	NP_076288.1
NCBI GenBank Nucleotide #	NM_023799.3
UniProt Accession #	Q9EQQ9; Q3JULY7; Q6ZQ71; Q8BK05; Q8BTT2; Q8CFX2; Q9CSJ4; Q9CUR7
Molecular Weight	107,395 Da
NCBI Official Full Name	protein O-GlcNAcase
NCBI Official Synonym Full Names	meningioma expressed antigen 5 (hyaluronidase)
NCBI Official Symbol	Mgea5
NCBI Official Synonym Symbols	Hy5; OGA; Ncoat; AA408215; mKIAA0679; 2810009A20Rik; 4833427O07Rik; 5830447M11Rik
NCBI Protein Information	protein O-GlcNAcase
UniProt Protein Name	Protein O-GlcNAcase
UniProt Gene Name	Mgea5
UniProt Synonym Gene Names	Hexc; Kiaa0679; OGA

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-elisa-kits/n-acetyl-beta-d-glucosaminidase-nagase/66678>