

Human Bruton'S Tyrosine Kinase (Btk) ELISA Kit | Btk elisa kit

Human Bruton'S Tyrosine Kinase (Btk) ELISA Kit

CATALOG #
AAA66683

REACTIVITY
Human


PRODUCT SYNONYMS

Bruton'S Tyrosine Kinase (Btk); Human Bruton'S Tyrosine Kinase (Btk) ELISA Kit; Btk elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Btk. No significant cross-reactivity or interference between Btk and analogues was observed.
Sequence Length	659
Samples	Tissue homogenates, cell lysates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.059ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Btk were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Btk were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for Btk elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of Btk in human tissue homogenates, cell lysates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to Btk. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to Btk. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Btk, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of Btk in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557377
NCBI GeneID	695
NCBI Accession #	NP_000052.1
NCBI GenBank Nucleotide #	NM_000061.2
UniProt Accession #	Q06187; Q32ML5; B2RAW1
Molecular Weight	79,937 Da
NCBI Official Full Name	tyrosine-protein kinase BTK isoform 1
NCBI Official Synonym Full Names	Bruton tyrosine kinase
NCBI Official Symbol	BTK
NCBI Official Synonym Symbols	AT; ATK; BPK; XLA; IMD1; AGMX1; PSCTK1
NCBI Protein Information	tyrosine-protein kinase BTK
UniProt Protein Name	Tyrosine-protein kinase BTK
UniProt Gene Name	BTK
UniProt Synonym Gene Names	AGMX1; ATK; BPK; ATK; BPK

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.