

## Human Osteocalcin (OC) ELISA Kit | OC elisa kit

Human Osteocalcin (OC) ELISA Kit

CATALOG #  
**AAA66698**

REACTIVITY  
**Human**


### PRODUCT SYNONYMS

Osteocalcin (OC); Human Osteocalcin (OC) ELISA Kit; BGLAP; OT; BGP; Bone Gla Protein; Bone Gamma-Carboxyglutamate Protein; Gamma-carboxyglutamic acid-containing protein; OC elisa kit

### PRODUCT SPECIFICATIONS

|                         |                                                                                                                                                                                                            |
|-------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Reactivity              | Human                                                                                                                                                                                                      |
| Specificity             | This assay has high sensitivity and excellent specificity for detection of OC. No significant cross-reactivity or interference between OC and analogues was observed.                                      |
| Sequence Length         | 100                                                                                                                                                                                                        |
| Samples                 | Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.                                                                                                      |
| Assay Type              | Sandwich                                                                                                                                                                                                   |
| Detection Range         | 62.5-4000pg/mL                                                                                                                                                                                             |
| Sensitivity             | 22.7pg/mL                                                                                                                                                                                                  |
| Intra-assay Precision   | Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level OC were tested 20 times on one plate, respectively. Intra-Assay: CV<10%                                       |
| Inter-assay Precision   | Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level OC were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%     |
| Preparation and Storage | For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C. |

### IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

|                                              |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|----------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Related Product Information for OC elisa kit | <p>Intended Uses: This kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of OC in human serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.</p> <p>Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to OC. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to OC. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain OC, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of OC in the samples is then determined by comparing the O.D. of the samples to the standard curve.</p> |
|----------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

NCBI AND UNIPROT INFORMATION

|                                  |                                     |
|----------------------------------|-------------------------------------|
| NCBI GI #                        | 40316933                            |
| NCBI GeneID                      | 632                                 |
| NCBI Accession #                 | NP_954642.1                         |
| NCBI GenBank Nucleotide #        | NM_199173.5                         |
| UniProt Accession #              | P02818; Q5TCK6                      |
| Molecular Weight                 | 10,963 Da                           |
| NCBI Official Full Name          | osteocalcin preproprotein           |
| NCBI Official Synonym Full Names | bone gamma-carboxyglutamate protein |
| NCBI Official Symbol             | BGLAP                               |
| NCBI Official Synonym Symbols    | OC; BGP; OCN                        |
| NCBI Protein Information         | osteocalcin                         |
| UniProt Protein Name             | Osteocalcin                         |
| UniProt Gene Name                | BGLAP                               |
| UniProt Synonym Gene Names       | BGP                                 |

**For Research Use Only.** Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | [contact@aaabiotech.com](mailto:contact@aaabiotech.com)  
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/osteocalcin-oc/66698>