

Human Thyroxine Binding Globulin (TBG) ELISA Kit | TBG elisa kit

Human Thyroxine Binding Globulin (TBG) ELISA Kit

CATALOG #
AAA66940

REACTIVITY
Human


PRODUCT SYNONYMS

Thyroxine Binding Globulin (TBG); Human Thyroxine Binding Globulin (TBG) ELISA Kit; SERPINA7 Serpin Peptidase Inhibitor Clade A Member 7 T4-binding globulin; TBG elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of TBG. No significant cross-reactivity or interference between TBG and analogues was observed.
Sequence Length	415
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids
Assay Type	Quantitative Sandwich
Detection Range	0.781-50ng/mL
Sensitivity	< 0.33ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level TBG were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level TBG were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for TBG elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of TBG in human serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to TBG. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to TBG. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain TBG, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of TBG in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	205277441
NCBI GeneID	6906
NCBI Accession #	NP_000345.2
NCBI GenBank Nucleotide #	NM_000354.5
UniProt Accession #	P05543; D3DUX1
Molecular Weight	46,325 Da
NCBI Official Full Name	thyroxine-binding globulin
NCBI Official Synonym Full Names	serpin family A member 7
NCBI Official Symbol	SERPINA7
NCBI Official Synonym Symbols	TBG; TBGQTL
NCBI Protein Information	thyroxine-binding globulin
UniProt Protein Name	Thyroxine-binding globulin
UniProt Gene Name	SERPINA7
UniProt Synonym Gene Names	TBG

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/thyroxine-binding-globulin-tbg/66940>