

Human Platelet Derived Growth Factor Subunit A (PDGFA) ELISA Kit | PDGFA elisa kit

Human Platelet Derived Growth Factor Subunit A (PDGFA) ELISA Kit

CATALOG #
AAA66952

REACTIVITY
Human


PRODUCT SYNONYMS

Platelet Derived Growth Factor Subunit A (PDGFA); Human Platelet Derived Growth Factor Subunit A (PDGFA) ELISA Kit; PDGF-A; PDGF1; Platelet Derived Growth Factor Alpha Polypeptide; PDGFA elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of PDGFA. No significant cross-reactivity or interference between PDGFA and analogues was observed.
Sequence Length	211
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.063ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level PDGFA were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level PDGFA were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for PDGFA elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of PDGFA in human serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to PDGFA. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to PDGFA. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain PDGFA, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of PDGFA in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	77695917
NCBI GeneID	5154
NCBI Accession #	NP_002598.4
NCBI GenBank Nucleotide #	NM_002607.5
UniProt Accession #	P04085; B5BU73
Molecular Weight	22,253 Da
NCBI Official Full Name	platelet-derived growth factor subunit A isoform 1 preproprotein
NCBI Official Synonym Full Names	platelet derived growth factor subunit A
NCBI Official Symbol	PDGFA
NCBI Official Synonym Symbols	PDGF1; PDGF-A
NCBI Protein Information	platelet-derived growth factor subunit A
UniProt Protein Name	Platelet-derived growth factor subunit A
UniProt Gene Name	PDGFA
UniProt Synonym Gene Names	PDGF1; PDGF subunit A

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/platelet-derived-growth-factor-subunit-a-pdgfa/66952>