

Human Glucose Transporter 2 (GLUT2) ELISA Kit | GLUT2 elisa kit

Human Glucose Transporter 2 (GLUT2) ELISA Kit

CATALOG #
AAA67071

REACTIVITY
Human


PRODUCT SYNONYMS

Glucose Transporter 2 (GLUT2); Human Glucose Transporter 2 (GLUT2) ELISA Kit; SLC2A2; SLC2-A2; Solute carrier family 2; facilitated glucose transporter member 2; Glucose transporter type 2; liver; GLUT2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of GLUT2. No significant cross-reactivity or interference between GLUT2 and analogues was observed.
Sequence Length	524
Samples	Serum, plasma, tissue homogenates and other biological fluids.
Assay Type	Sandwich
Detection Range	1.56-100ng/mL
Sensitivity	0.64ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level GLUT2 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level GLUT2 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for GLUT2 elisa kit	Intended Uses: This kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of GLUT2 in human serum, plasma, tissue homogenates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to GLUT2. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to GLUT2. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain GLUT2, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of GLUT2 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557851
NCBI GeneID	6514
NCBI Accession #	NP_000331.1
NCBI GenBank Nucleotide #	NM_000340.1
UniProt Accession #	P11168; Q9UCW9; A8K481; B2R936; B7Z547; F8W8V8
Molecular Weight	44,702 Da
NCBI Official Full Name	solute carrier family 2, facilitated glucose transporter member 2 isoform 1
NCBI Official Synonym Full Names	solute carrier family 2 member 2
NCBI Official Symbol	SLC2A2
NCBI Official Synonym Symbols	GLUT2
NCBI Protein Information	solute carrier family 2, facilitated glucose transporter member 2
UniProt Protein Name	Solute carrier family 2, facilitated glucose transporter member 2
UniProt Gene Name	SLC2A2
UniProt Synonym Gene Names	GLUT2; GLUT-2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.