

Mouse DNA Damage Inducible Transcript 3 (DDIT3) ELISA Kit | DDIT3 elisa kit

Mouse DNA Damage Inducible Transcript 3 (DDIT3) ELISA Kit

CATALOG #
AAA67101

REACTIVITY
Mouse


PRODUCT SYNONYMS

DNA Damage Inducible Transcript 3 (DDIT3); Mouse DNA Damage Inducible Transcript 3 (DDIT3) ELISA Kit; DDIT3 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of DDIT3. No significant cross-reactivity or interference between DDIT3 and analogues was observed.
Sequence Length	168
Samples	Tissue homogenates, cell lysates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.058ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level DDIT3 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level DDIT3 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for DDIT3 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of DDIT3 in mouse tissue homogenates, cell lysates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to DDIT3. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to DDIT3. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain DDIT3, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of DDIT3 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	589058168
NCBI GeneID	13198
NCBI Accession #	NP_001277112.1
NCBI GenBank Nucleotide #	NM_001290183.1
UniProt Accession #	P35639; Q91YW9
Molecular Weight	19,189 Da
NCBI Official Full Name	DNA damage-inducible transcript 3 protein
NCBI Official Synonym Full Names	DNA-damage inducible transcript 3
NCBI Official Symbol	Ddit3
NCBI Official Synonym Symbols	chop; CHOP10; CHOP-10; gadd153
NCBI Protein Information	DNA damage-inducible transcript 3 protein
UniProt Protein Name	DNA damage-inducible transcript 3 protein
UniProt Gene Name	Ddit3
UniProt Synonym Gene Names	Chop; Chop10; Gadd153; DDIT-3; CHOP; CHOP-10

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-elisa-kits/dna-damage-inducible-transcript-3-ddit3/67101>