

Human Matrix Gla Protein (MGP) ELISA Kit | MGP elisa kit

Human Matrix Gla Protein (MGP) ELISA Kit

CATALOG #
AAA67121

REACTIVITY
Human


PRODUCT SYNONYMS

Matrix Gla Protein (MGP); Human Matrix Gla Protein (MGP) ELISA Kit; IL2R-a; CD25; IL2-RA; IDDM10; IL2R; TCGFR; MGP elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of MGP. No significant cross-reactivity or interference between MGP and analogues was observed.
Sequence Length	103
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	15.625-1000pg/mL
Sensitivity	6.7pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level MGP were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level MGP were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for MGP elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of MGP in human serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to MGP. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to MGP. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain MGP, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of MGP in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	49574514
NCBI GeneID	4256
NCBI Accession #	NP_000891.2
NCBI GenBank Nucleotide #	NM_000900.4
UniProt Accession #	P08493; Q2TU41; Q567P9; Q6ICN5; A0M8W5; B2R519; J3KMX7
Molecular Weight	15,325 Da
NCBI Official Full Name	matrix Gla protein isoform 2 preproprotein
NCBI Official Synonym Full Names	matrix Gla protein
NCBI Official Symbol	MGP
NCBI Official Synonym Symbols	NTI; GIG36; MGLAP
NCBI Protein Information	matrix Gla protein
UniProt Protein Name	Matrix Gla protein
UniProt Gene Name	MGP
UniProt Synonym Gene Names	MGLAP; MGP

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.